Research Article

The comparison of cosmetic results of Karydakis and Limberg flap Techniques in pilonidal sinus

Umit Sekmen^{1,3®}, Hamit Karayagiz^{2®}, Melih Paksoy^{3®}

¹Fulya Academy Clinic, General Surgery, Istanbul, Türkiye ²General Surgery, Esenyurt Necmi Kadioglu State Hospital, Istanbul, Türkiye ³General Surgery, Acıbadem Fulya Hospital, Istanbul, Türkiye

ABSTRACT

Background: Repairing of pilonidal sinus with flap techniques is very effective in prevention of recurrence. Karydakis and Limberg are two popular flap techniques. Their postoperative clinical outcomes are almost the same in the literature. Recently, all surgical techniques try to be less invasive as minimal as possible with the best cosmetic outcome. The aim of this study is to compare these two techniques in terms of cosmetic results.

Methods: Total 100 patients operated with Karydakis or Limberg were devided into two groups for comparison of their cosmetic results. Age, gender, BMI and postoperative period are used for demographic comparison and Stony Brook Scar Evaluation Scale (SBSES) was used to compare the cosmetic results of both surgical flap techniques.

Results: We found no difference between the groups in terms of demographic features but the SBSES scores was found higher in Karydakis group.

Conclusion: Karydakis technique might be preffered more than limberg technique with its better cosmetic result.

Keywords: minimal invasive, cosmetic, pilonidal sinus, flap technique

Introduction

Pilonidal sinus is the most common chronic disease of the sacrococcygeal region. Even most of the studies reported an incidence of 26/100,000 in the worldwide, in some of them the incidence was found around 8% among men, that is more common especially in Mediterranean region countries (1-5). It was first described by Herbert Mayo in 1833. Treatment options range from microsinusectomy to excision with flap repairing (6). Karydakis and

limberg flap repairs are most popular surgery techniques in the literature, and also mostly preferred techniques in our clinic. The search for minimally invasive and more cosmetic techniques with similar clinical results must be the target of the preferred treatment technique. No studies have been conducted comparing these two techniques only in terms of objective cosmetic results. We aimed to compare the cosmetic results of these two techniques, those have almost the same clinical results in the literature.

[⊠] Umit Sekmen • usekmen@yahoo.com

Received: 23.11.2024 • Accepted 11.03.2025

Copyright © 2025 The Author(s). This is an open access article distributed under the Creative Commons Attribution License (CC BY), which permits unrestricted use, distribution, and reproduction in any medium or format, provided the original work is properly cited.

Patients and Methods

Patients, who underwent karydakis and limberg flap repair surgery for pilonidal sinus disease in Acıbadem Fulya Hospital and Esenyurt State Hospital general surgery clinics in between 2019 and 2023, were invited by phone for clinical examination. First 50 patients for each surgery group who accepted to be in our study were included in this study. Patients with recurrence or who underwent surgical intervention due to recurrence were not included in the study. Two separate groups were formed as Karydakis group (n=50) and Limberg group (n=50). Both groups were evaluated according to demographic features; age, gender and body mass index (BMI).

For comparison of cosmetic results of these two techniques, Stony Brook Scar Evaluation Scale (SBSES), a validated scale specifically to

Table 1. Stony Brook Scar Evaluation Scale (SBSES)

I Trends Med Invest 2025;1(1):15-19

measure the long-term appearance of scars was used (Table 1) (7). For statistical analysis SPSS Student -t test was used.

The study was designed in accordance with Principles of the Declaration of Helsinki and ethically approved by Committee of Ethics of Acıbadem University (2020/05).

Results

The mean age was 24 (17-38), and the femalemale ratio was 16/84. There was no difference between the groups in terms of postoperative period, gender ratio, and BMI. The cosmetic scores of the karydakis group were found to be significantly higher. Demographic features of the patients are summarized in Table 2, and SBSES scores are summarized in Table 3.

	Scar Category	Points
Width	>2 mm	0
	< 2 mm	1
Height	Elevated/Depressed in relation to surrounding skin	0
	Flat	1
Color	Darker than surrounding skin	0
	Same color or lighter than surrounding skin	1
Suture Marks	Present	0
	Absent	1
Overall Appearance	Poor	0
	Good	1

Total Score = Sum of individual score, from 0 (worst score) to 5 (best score)

Table 2. Demographics of Patients

	Group 1	Group 2	Р
Age (Mean)	26.45±5.35	27.73±6.18	0.08
F/M Ratio	9/41 (21.9%)	7/43 (16.3%)	0.07
BMI	24.53±5.5	25.12±5.8	0.1
Mean	15.5±3.3	16.2±3.6	0.1
Postoperative			
Period (Month)			

Table 3. SBSES Scores

	Group 1	Group 2	Р
Width	43	32	0.02
Height	46	29	0.01
Color	47	48	0.09
Suture Marks	14	9	0.01
Overall Appearence	47	36	0.04
Mean Total Score	39.4	30.8	0.01

Discussion

Even though there are some studies still searching whether pilonidal sinus disease is a congenital or acquired disease, the common view is that it is more of an acquired disease (8-10). The treatment of the disease is surgery, and lower recurrence rates make surgeries performed with flaps advantageous over primary repair, making it the preferred surgical technique (11-15). Comparing these 2 most preferred flap techniques (Karydakis and Limberg flaps) in their systematic review and meta-analysis, Gavriilidis et al. evaluated both techniques in terms of recurrence and complications and concluded that there was no significant difference between them (16).

With the same clinical results, one of these two techniques is preferred according to surgeon's choice. Although these techniques were slightly superior to each other in a limited number of studies comparing operative time, early and late postoperative complications, and recurrence rates, they were not superior to each other in systematic reviews and metaanalyses conducted on this subject (16-18). In the study of Erkent et al., they stated that in the retrospective analysis of 924 cases with flap and primary closure, the primary closure technique was preferred especially in female patients, and cosmetic preferences were at the forefront in this (19).

In the literature, satisfaction questions regarding the incisions were asked to patients after the surgery, while similar rates were reported in the studies, some studies reported that they found a statistically higher satisfaction rate for Karydakis (20-24). However, in a few of these clinical studies, patients were questioned in terms of scar satisfaction, and it was found higher in Karydakis group (25,26).

Our study is original in terms of more objective comparison of cosmetic results. The scale, which include the evaluations of both the patient and the surgeon, is the first objective scale using in a study to compare these two techniques cosmetically. According to SBSES, our study found the cosmetic score to be significantly higher in the Karydakis group than in the Limberg group.

So we may conclude that Karydakis technique may be preffered instead of Limberg technique with its better cosmetic outcome.

Conclusion

Pilonidal sinus is a common disease of the intergluteal region. Flap repairs are very effective in its treatment. Karydakis and Limberg flap techniques are the 2 important surgical techniques preferred in this region diseases. Today, where minimally invasive and more cosmetic surgery options are more preferred, we aimed to compare these two surgical techniques in terms of cosmetic outcome. For such an objective study, which has never been done before, we used the Stony Brook Scar Evaluation Scale. As a result of our study, we found the cosmetic score of the Karydakis flap technique to be higher than the limberg flap technique.

As a result, we revealed out that the choice of the karydakis flap in intergluteal pilonidal sinus disease has better cosmetic results.

Ethical approval

The study was approved by Acıbadem University Ethics Committee (2020/05).

Author contribution

The authors confirm contribution to the paper as follows: Study conception and design: US; data collection: HK; analysis and interpretation of results: HK; draft manuscript preparation: US, MP. All authors reviewed the results and approved the final version of the article. Sekmen U, et al.

Source of funding

The authors declare the study received no funding.

Conflict of interest

The authors declare that there is no conflict of interest.

REFERENCES

- Sekmen U, Kara VM, Altintoprak F, Senol S. Pilonidal sinus in the army: Its incidence and risk factors. Turk J Surg. 2010;26:95-8. [Crossref]
- Søndenaa K, Andersen E, Nesvik I, Søreide JA. Patient characteristics and symptoms in chronic pilonidal sinus disease. Int J Colorectal Dis. 1995;10(1):39-42. [Crossref]
- 3. Al-Khamis A, McCallum I, King PM, Bruce J. Healing by primary versus secondary intention after surgical treatment for pilonidal sinus. Cochrane Database Syst Rev. 2010;2010(1):CD006213. [Crossref]
- Akinci OF, Kurt M, Terzi A, Atak I, Subasi IE, Akbilgic O. Natal cleft deeper in patients with pilonidal sinus: implications for choice of surgical procedure. Dis Colon Rectum. 2009;52(5):1000-2. [Crossref]
- Lee PJ, Raniga S, Biyani DK, Watson AJ, Faragher IG, Frizelle FA. Sacrococcygeal pilonidal disease. Colorectal Dis. 2008;10(7):639-50; discussion 651-2. [Crossref]
- 6. Mayo OH. Observations on injuries and diseases of rectum. London: Burgess and Hill; 1833: 45-6.
- Singer AJ, Arora B, Dagum A, Valentine S, Hollander JE. Development and validation of a novel scar evaluation scale. Plast Reconstr Surg. 2007;120(7):1892-1897. [Crossref]
- Hodges RM. Pilo-nidal sinus. Boston Med Surg J. 1880;103(21):485-6. [Crossref]
- Bascom J, Bascom T. Failed pilonidal surgery: new paradigm and new operation leading to cures. Arch Surg. 2002;137(10):1146-50; discussion 1151. [Crossref]
- Karydakis GE. New approach to the problem of pilonidal sinus. Lancet. 1973;2(7843):1414-5. [Crossref]

- Akca T, Colak T, Ustunsoy B, Kanik A, Aydin S. Randomized clinical trial comparing primary closure with the Limberg flap in the treatment of primary sacrococcygeal pilonidal disease. Br J Surg. 2005;92(9):1081-4. [Crossref]
- Can MF, Sevinc MM, Yilmaz M. Comparison of Karydakis flap reconstruction versus primary midline closure in sacrococcygeal pilonidal disease: results of 200 military service members. Surg Today. 2009;39(7):580-6. [Crossref]
- Arslan K, Said Kokcam S, Koksal H, Turan E, Atay A, Dogru O. Which flap method should be preferred for the treatment of pilonidal sinus? A prospective randomized study. Tech Coloproctol. 2014;18(1):29-37. [Crossref]
- Keshvari A, Keramati MR, Fazeli MS, Kazemeini A, Meysamie A, Nouritaromlou MK. Karydakis flap versus excision-only technique in pilonidal disease. J Surg Res. 2015;198(1):260-6. [Crossref]
- Arnous M, Elgendy H, Thabet W, Emile SH, Elbaz SA, Khafagy W. Excision with primary midline closure compared with Limberg flap in the treatment of sacrococcygeal pilonidal disease: a randomised clinical trial. Ann R Coll Surg Engl. 2019;101(1):21-9. [Crossref]
- Gavriilidis P, Bota E. Limberg flap versus Karydakis flap for treating pilonidal sinus disease: a systematic review and meta-analysis. Can J Surg. 2019;62(2):131-8. [Crossref]
- 17. Bessa SS. Comparison of short-term results between the modified Karydakis flap and the modified Limberg flap in the management of pilonidal sinus disease: a randomized controlled study. Dis Colon Rectum. 2013;56(4):491-8. [Crossref]
- Sahebally SM, McMahon G, Walsh SR, Burke JP. Classical Limberg versus classical Karydakis flaps for pilonidal disease- an updated systematic review and meta-analysis of randomized controlled trials. Surgeon. 2019;17(5):300-8. [Crossref]
- Erkent M, Şahiner İT, Bala M, et al. Comparison of Primary Midline Closure, Limberg Flap, and Karydakis Flap Techniques in Pilonidal Sinus Surgery. Med Sci Monit. 2018;24:8959-63. [Crossref]
- Ates M, Dirican A, Sarac M, Aslan A, Colak C. Short and long-term results of the Karydakis flap versus the Limberg flap for treating pilonidal sinus disease: a prospective randomized study. Am J Surg. 2011;202(5):568-73. [Crossref]

J Trends Med Invest 2025;1(1):15-19

- 21. Tokac M, Dumlu EG, Aydin MS, Yalcın A, Kilic M. Comparison of modified Limberg flap and Karydakis flap operations in pilonidal sinus surgery: prospective randomized study. Int Surg. 2015;100(5):870-7. [Crossref]
- 22. Bali İ, Aziret M, Sözen S, et al. Effectiveness of Limberg and Karydakis flap in recurrent pilonidal sinus disease. Clinics (Sao Paulo). 2015;70(5):350-5. [Crossref]
- 23. Khan KJ, Ghaffar A, Choudhry S, Irshad K. Comparison of early outcome between modified Limberg and Karydakis flap procedures in patients with sacrococcygeal pilonidal sinus. Pak J Med Health Sci. 2016;10:631-4.
- 24. Milone M, Di Minno MND, Musella M, et al. The role of drainage after excision and primary closure of pilonidal sinus: a meta-analysis. Tech Coloproctol. 2013;17(6):625-30. [Crossref]
- 25. Eryilmaz R, Sahin M, Alimoglu O, Dasiran F. Surgical treatment of sacrococcygeal pilonidal sinus with the Limberg transposition flap. Surgery. 2003;134(5):745-9. [Crossref]
- 26. Yildiz MK, Ozkan E, Odabaşı HM, et al. Karydakis flap procedure in patients with sacrococcygeal pilonidal sinus disease: experience of a single centre in Istanbul. ScientificWorldJournal. 2013;2013:807027. [Crossref]